



Approaches to Managing Cardiovascular Risk Factors in Patients with Type 2 Diabetes Mellitus

Mokhmad-Emi Turkulovich Ilyasov¹, Madina Valerievna Nadgerieva², Magomed Adamovich Dautmerzaev³, Alibek Muratovich Baitemirov⁴, Daria Alexandrovna Zhezheria⁵, Yuriy Borisovich Fedoseyev⁵, Angelina Georgievna Shevchenko¹, Anastasia Stanislavovna Khutatova⁶, Kavsarat Kamaldinovna Atagishieva⁴, Zhamilat Arsenovna Abdullaeva⁴, Elena Valentinovna Svetlakova⁷

¹ Stavropol State Medical University, Stavropol, RUSSIA.

² North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, RUSSIA.

³ Saratov State Medical University named after Razumovsky, Saratov, RUSSIA.

⁴ Dagestan State Medical University, Makhachkala, Republic of Dagestan, RUSSIA.

⁵ Rostov State Medical University, Rostov-on-Don, RUSSIA.

⁶ Nevinnomyssk City Children's Hospital, Stavropol Region, RUSSIA.

⁷ Stavropol State Agrarian University, Stavropol, RUSSIA.

*Corresponding Author (Tel: +79183500889, Email: ruslankalmykov777@yandex.ru).

Paper ID: 13A11A

Volume 13 Issue 11

Received 23 July 2021

Received in revised form 19 October 2022

Accepted 26 October 2022

Available online 01 November 2022

Keywords:

Type 2 diabetes mellitus;
Cardiovascular diseases;
Risk factors;
Cardiovascular risk management.

Abstract

Diabetes mellitus is one of the most common diseases, and therefore, this problem is one of the priorities for doctors and healthcare in all countries. Type 2 diabetes mellitus is a chronic disease that leads to the risk of developing cardiovascular complications and pathologies in patients, and, as a result, to high mortality or disability. Diabetes is combined with such diseases as coronary heart disease, stroke, atherosclerosis of the vessels of the lower extremities, cardiomyopathy, and congestive heart failure. This article reviews the treatment management of diabetes, especially to reduce mortality among patients with diabetes mellitus.

Discipline: Medicine.

©2022 INT TRANS J ENG MANAG SCI TECH.

Cite This Article:

Ilyasov, M.-E.T., Nadgerieva, M.V., Dautmerzaev, M.A., Baitemirov, A.M., Zhezheria, D.A., Fedoseyev, Yu.B., ... Svetlakova, E.V. (2022). Approaches to Managing Cardiovascular Risk Factors in Patients with Type 2 Diabetes Mellitus. *International Transaction Journal of Engineering, Management, & Applied Sciences & Technologies*, 13(11), 13A11A, 1-7. <http://TUENGR.COM/V13/13A11A.pdf> DOI: 10.14456/ITJEMAST.2022.211

1 Introduction

The problem of diabetes mellitus has high growth and prevalence everywhere from year to year, and therefore its relevance is increasing more and more [1].

Weight gain, the development of metabolic syndrome, and type 2 diabetes mellitus may occur as a response to:

- lack of movement (hypokinesia);
- improper nutrition: high content of fats, salts, carbohydrates, and various additives in food;
- bad habits;
- stress factors;
- acceleration of the pace of life;
- genetics;
- ecology.

Thus, at this stage, diabetes mellitus, as a non-communicable disease, is on par with oncology and cardiovascular pathology, leading to disability or death.

2 The Current State of the Issue of Diabetes Mellitus

Diabetes mellitus refers to chronic diseases and has a progressive type, is characterized by an increased level of glucose in the blood serum (hyperglycemia), and has concomitant conditions (deterioration of vision, poor wound healing, kidney failure, cardiovascular diseases, etc.) [2].

Of particular concern is the fact that in 50% of cases the disease is not diagnosed [3,4]. Often, the disease can be detected only a few years after its occurrence, which leads to the presence of complications.

Researchers have found that a large number of patients with diabetes have type 2 diabetes mellitus. Type 2 diabetes mellitus is adult diabetes or insulin-dependent. The causes of its occurrence are an unhealthy lifestyle and insufficient physical activity. Until recently, such diabetes was detected only in adults, but the disease has received a "rejuvenation" and can now be detected in children [5].

The development of diabetes occurs in cases of insufficient production and effective use of insulin. With this disease, the level of sugar in the blood increases, and as a result, this leads to malfunctions in the body's system. It has a special effect on the nervous and cardiovascular systems [6-8].

The spread of diabetes mellitus depends on the level of income of the population: the lower it is, the faster the disease develops. In low-income countries, a particular prevalence of diabetes is observed in people over 60 years old, and in high-income countries - in patients after 70-75 years.

3 Cardiovascular Risk Factors in Patients with Type 2 Diabetes Mellitus and Approaches to Their Management

One of the main causes of death in patients with diabetes mellitus is cardiovascular pathology, and diabetes mellitus itself is a cardiovascular risk factor. In addition, diabetes is

combined with such diseases as coronary heart disease, stroke, atherosclerosis of the vessels of the lower extremities, cardiomyopathy, and congestive heart failure [9-11].

With type 2 diabetes, metabolic disorders appear that form risk factors for the cardiovascular system (obesity, dyslipidemia, hyperinsulinemia, insulin resistance, etc.).

The relationship between elevated blood sugar (hyperglycemia) and cardiovascular diseases is very high. With chronic hyperglycemia, atherosclerosis often develops (free radicals are intensively formed, which can bind to lipid molecules) [12,13]. An increase in the level of glycemia increases the risk of developing cardiovascular pathology.

The indicator reflecting the average daily plasma glucose level is glycated hemoglobin (HbA1c). The HbA1c level has a standard and determines the glycemic status in patients with diabetes mellitus.

Scientists have also revealed that complications of diabetes mellitus can occur from the variability of glycemia. With its strong increase, powerful oxidative stress is observed. Patients with type 2 diabetes mellitus are more likely to develop ventricular arrhythmias if there is a high amplitude of daily fluctuations in glycemia [14].

Also, fluctuations in glycemia can lead to disruption of vascular processes (for example, cause inflammation or oxidative stress). Such fluctuations most often occur in patients with type 2 diabetes mellitus and depend on meals.

That is why strict glycemic control is recommended, which will minimize the risk of complications, as well as reduce the mortality rate [15]. But at the same time, there may be negative results: weight gain, hypoglycemia, etc. Thus, the choice of medications and the selection of therapy are carried out individually, taking into account the characteristics of each patient's body.

Insulin resistance (IR) is one of the causes of the occurrence and rapid development of cardiovascular diseases. At the same time, the frequency of multiple atherosclerotic lesions of coronary vessels increases. Insulin resistance adversely affects the structure and function of the myocardium in patients of different risk categories.

Another factor influencing cardiovascular diseases is obesity [16-19]. Adipokines such as leptin, adiponectin, resistin, omentin, visfatin, etc. are produced in adipose tissue. Leptin, namely its level positively interacts with the content of triglycerides and free fatty acids in patients with type 2 diabetes mellitus and obesity, which indicates the relationship of leptin with lipotoxicity [20].

To reduce overall mortality, it is necessary to reduce body weight, and at the same time, this will have a positive effect on glycemic control in patients with type 2 diabetes [21-23].

Hyperuricemia - an increased content of uric acid also serves as a risk factor for the development of cardiovascular diseases. Elevated uric acid levels can lead to arterial hypertension, congestive heart failure, and impaired kidney function [24,25].

For patients with cardiovascular risk, the elevated uric acid content is especially dangerous and can be a prognostic parameter of cardiovascular and general mortality.

Thus, disorders of uric acid metabolism in patients with diabetes mellitus should be carefully monitored and corrected in time.

To reduce mortality among patients with diabetes mellitus, it is necessary to control and manage cardiovascular risk factors [26, 27]. This is possible thanks to the use of individually selected drugs that have favorable properties for the cardiovascular system.

A hypoglycemic drug selected by a specialist should perform several functions at once: firstly, to control glycemia, and secondly, to minimize the risk of developing cardiovascular diseases.

Particular attention is paid to the effect of hypoglycemic drugs on reducing the volume of visceral fat. Such fat causes complications in type 2 diabetes mellitus, and the use of drugs reduces its volume [28,29].

Also, for many patients, hypoglycemic therapy is recommended, which will be of a combined nature [30].

4 Conclusion

Type 2 diabetes mellitus is a very common disease that is of particular importance for all countries. Cardiovascular risks pose a particular threat to the life of patients with type 2 diabetes mellitus (according to statistics, the mortality rate is above 50%). That is why such a strategy of treatment of patients with type 2 diabetes mellitus is recommended, in which there will be a one-time impact on cardiovascular risk factors.

Researchers recommend developing an individual treatment line in which the assessment of hypoglycemic drugs would be considered not only because of their effect on the compensation of carbohydrate metabolism but also took into account the cardiovascular risks of patients and the possibility of their further correction. At the same time, it is necessary to pay attention to cardiovascular diseases that have already been detected in the patient. Thus, when choosing a drug, it is necessary to conduct studies and analyzes in order to exclude a negative impact on the course of the disease.

5 Availability of Data and Material

Data can be made available by contacting the corresponding author.

6 References

- [1] Gerstein HC, Colhoun HM, Dagenais GR, Diaz R, Lakshmanan M, Pais P, Probstfield J, Riesenmeyer JS, Riddle MC, Rydén L, Xavier D. Dulaglutide and cardiovascular outcomes in type 2 diabetes (REWIND): a double-blind, randomised placebo-controlled trial. *The Lancet*. 2019 Jul 13;394(10193):121-30.
- [2] González-Lao E, Corte Z, Simón M, Ricós C, Coskun AB, Braga F, Aarsand AK, Carobene A, Bartlett WA, Boned B, Asland B. Systematic review of the biological variation data for diabetes related analytes. *Clinica Chimica Acta*. 2019 Jan 1;488:61-7.
- [3] Raevskaya AI, Belyalova AA, Shevchenko PP, Karpov SM, Mishvelov AE, Simonov AN, Povetkin SN. Cognitive Impairments in a Range of Somatic Diseases. *Diagnostics, Modern Approach to Therapy. People*;22:23.
- [4] Bledzhyants GA, Mishvelov AE, Nuzhnaya KV, Anfinogenova OI, Isakova JA, Melkonyan RS, Hite GY,

- Suprunchuk VE, Makova AV, Popov AN, Ovechkin VS. The effectiveness of the medical decision-making support system" electronic clinical pharmacist" in the management of patients therapeutic profile. Artificial intelligence. 2019 Apr 28.
- [5] Boraeva TT, Vadaeva MA, Matveeva UV, Revazova AB, Albegova BZ, Kanukoeva DT, Mishvelov AE, Povetkin SN. Dynamics of Diseases of the Upper Digestive Tract in Children. *Journal of Pharmaceutical Research International*. 2021;33(38B):48-57.
- [6] Michos ED, McEvoy JW, Blumenthal RS. Lipid management for the prevention of atherosclerotic cardiovascular disease. *New England Journal of Medicine*. 2019 Oct 17;381(16):1557-67.
- [7] Grenier J, Goodman SG, Leiter LA, Langer A, Teoh H, Bhatt DL, Cheng AY, Tan MK, Ransom T, Connelly KA, Yan AT. Blood pressure management in adults with type 2 diabetes: insights from the diabetes mellitus status in Canada (DM-SCAN) survey. *Canadian Journal of Diabetes*. 2018 Apr 1;42(2):130-7.
- [8] Siddiqui SA, Ali Redha A, Snoeck ER, Singh S, Simal-Gandara J, Ibrahim SA, Jafari SM. Anti-Depressant Properties of Crocin Molecules in Saffron. *Molecules*. 2022 Mar 23;27(7):2076.
- [9] Giorgino F, Vora J, Fenici P, Solini A. Renoprotection with SGLT2 inhibitors in type 2 diabetes over a spectrum of cardiovascular and renal risk. *Cardiovascular Diabetology*. 2020 Dec;19(1):1-9.
- [10] Johnson RJ, Bakris GL, Borghi C, Chonchol MB, Feldman D, Lanasa MA, Merriman TR, Moe OW, Mount DB, Lozada LG, Stahl E. Hyperuricemia, acute and chronic kidney disease, hypertension, and cardiovascular disease: report of a scientific workshop organized by the National Kidney Foundation. *American Journal of Kidney Diseases*. 2018 Jun 1;71(6):851-65.
- [11] Mezhidov, B. S., Bekova, K. I., Edieva, A. A., Madalova, K. I., Malakhova, V. A., Uspaeva, Z. A., Tachaeva, M. S., Tsinaeva, A. M., Maslova, A. Y. and Khodisov, S. V. (2021) "Association of Genetic Polymorphism of Tumor Necrosis Factor- α in the Development of Coronary Heart Disease in Elderly Patients", *Journal of Pharmaceutical Research International*, 33(54B), pp. 81-88. doi: 10.9734/jpri/2021/v33i54B33768
- [12] Selimov MA, Nagdalian AA, Povetkin SN, Statsenko EN, Kulumbekova IR, Kulumbekov GR, Litvinov MS, Mishvelov AE, Iraskhanov AS, Abdurashidov AM, Gadzhiev MR. Investigation of CdCl₂ Influence on Red Blood Cell Morphology. *International Journal of Pharmaceutical and Phytopharmacological Research*. 2019;9(5):8-13.
- [13] Rzhepakovsky I, Anusha Siddiqui S, Avanesyan S, Benlidayi M, Dhingra K, Dolgalev A, Enuakashvily N, Fritsch T, Heinz V, Kochergin S, Nagdalian A. Anti-arthritic effect of chicken embryo tissue hydrolyzate against adjuvant arthritis in rats (X-ray microtomographic and histopathological analysis). *Food Science & Nutrition*. 2021 Oct;9(10):5648-69.
- [14] Ungurianu A, Zanzfirescu A, Grădinaru D, Ionescu-Tîrgoviște C, Dănciulescu Miulescu R, Margină D. Interleukins and redox impairment in type 2 diabetes mellitus: mini-review and pilot study. *Current Medical Research and Opinion*. 2022 Apr 3;38(4):511-22.
- [15] Viigimaa M, Sachinidis A, Toumpourleka M, Koutsampasopoulos K, Alliksoo S, Titma T. Macrovascular complications of type 2 diabetes mellitus. *Current vascular pharmacology*. 2020 Mar 1;18(2):110-6.
- [16] Rybak V, Derkach N, Korol V. Modern aspects of sugar-reducing effect of thick bean extract based on a type ii diabetes model on the background of obesity. *Georgian medical news*. 2019 May 1;5(290):144-9.
- [17] Dong N, Gao YH, Liu B, Zhao CW, Yang C, Li SQ, Liu JG, Qi X. Differential expression of adipokines in knee osteoarthritis patients with and without metabolic syndrome. *International Orthopaedics*. 2018 Jun;42(6):1283-9.
- [18] Ayivi R, Ibrahim SA, Colleran H, Silva R, Williams L, Galanakis C, Fidan H, Tomovska J, Siddiqui SA. COVID-19: human immune response and the influence of food ingredients and active compounds. *Bioactive Compounds in Health and Disease*. 2021 Jun 29;4(6):100-48.
- [19] Ranjha MM, Shafique B, Rehman A, Mehmood A, Ali A, Zahra SM, Nasreen S, Singh A, Ibrahim SA, Siddiqui SA. Biocompatible Nanomaterials in Food Science, Technology and Nutrient Drug Delivery: Recent

Developments and Applications. *Frontiers in Nutrition*. 2021 Jan 1:1141.

- [20] Sergeevna LM, Ivanovich KV, Sergey P, Viktorovich PS, MarinaPetrovna M, Viktorovich SV, Ashotovich76 NA. 5% Suspension of albendazole echinacea magenta (*echinacea purpurea*) toxicometric evaluation. *Entomology and Applied Science Letters*. 2018 Jan 1;5(4):30-4.
- [21] Churuangsuk C, Hall J, Reynolds A, Griffin SJ, Combet E, Lean ME. Diets for weight management in adults with type 2 diabetes: an umbrella review of published meta-analyses and systematic review of trials of diets for diabetes remission. *Diabetologia*. 2021 Nov 17:1-23.
- [22] Maggio CA, Pi-Sunyer FX. Obesity and type 2 diabetes. *Endocrinology and Metabolism Clinics*. 2003 Dec 1;32(4):805-22.
- [23] Lavie CJ, Milani RV, Ventura HO. Obesity and cardiovascular disease: risk factor, paradox, and impact of weight loss. *Journal of the American college of cardiology*. 2009 May 26;53(21):1925-32.
- [24] Katsiki N, Papanas N, Fonseca VA, Maltezos E, Mikhailidis DP. Uric acid and diabetes: is there a link?. *Current pharmaceutical design*. 2013 Aug 1;19(27):4930-7.
- [25] Areshidze DA, Mischenko DV, Makartseva LA, Kucher SA, Kozlova MA, Timchenko LD, Rzhepakovsky IV, Nagdalian AA, Pushkin SV. Some functional measures of the organism of rats at modeling of ischemic heart disease in two different ways. *Entomology and Applied Science Letters*. 2018 Jan 1;5(4):19-29.
- [26] Lai YJ, Chen YY, Ku PW, Chen LJ, Yen YF. Association between uric acid level and incidence of albuminuria in patients with type 2 diabetes mellitus: A 4.5-year cohort study. *Medicine*. 2021 Oct 10;100(41).
- [27] Lou Y, Qin P, Wang C, Ma J, Peng X, Xu S, Chen H, Zhao D, Wang L, Liu D, Li Y. Sex-specific association of serum uric acid level and change in hyperuricemia status with risk of type 2 diabetes mellitus: a large cohort study in China. *Journal of Diabetes Research*. 2020 Jul 24;2020.
- [28] Bloomgarden Z. Diabetes and branched-chain amino acids: What is the link?. *Journal of diabetes*. 2018 Feb 13;10(5):350-2.
- [29] Kim HS, Shin JA, Lee SH, Kim ES, Cho JH, Son HY, Yoon KH. A comparative study of the effects of a dipeptidyl peptidase-IV inhibitor and sulfonylurea on glucose variability in patients with type 2 diabetes with inadequate glycemic control on metformin. *Diabetes technology & therapeutics*. 2013 Oct 1;15(10):810-6.
- [30] Vinik AI, Nevoret ML, Casellini C, Parson H. Diabetic neuropathy. *Endocrinology and Metabolism Clinics*. 2013 Dec 1;42(4):747-87.
-



Mokhmad-Emi Turkulovich Ilyasov is a student of Stavropol State Medical University, Stavropol, Russia



Madina Valerievna Nadgerieva is a student of the North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Magomed Adamovich Dautmerzaev is a student of Saratov State Medical University named after Razumovsky, Saratov, Russia



Alibek Muratovich Baitemirov is a student of Dagestan State Medical University, Makhachkala, Republic of Dagestan, Russia




Daria Alexandrovna Zhezheria is a student of Rostov State Medical University, Rostov-on-Don, Russia



Yuriy Borisovich Fedoseyev is a student of Rostov State Medical University, Rostov-on-Don, Russia



Angelina Georgievna Shevchenko is a student of Stavropol State Medical University, Stavropol, Russia



Anastasia Stanislavovna Khutatova, Nevinnomyssk City Children's Hospital, Stavropol Region, Russia



Kavsarat Kamaldinovna Atagishieva is a student of Dagestan State Medical University, Makhachkala, Republic of Dagestan, Russia



Zhamilat Arsenovna Abdullaeva is a student of Dagestan State Medical University, Makhachkala, Republic of Dagestan, Russia



Dr. Elena Valentinovna Svetlakova is an associate professor, Department of epizootology and microbiology, Faculty of Veterinary Medicine, Stavropol State Agrarian University, Stavropol, Russia