



Possibilities of Correction of Functional Diseases of the Gastrointestinal Tract in Patients with Anxiety Disorders

Maksalina Ilyasovna Makaeva¹, Glikeria Levanovna Khabalaeva¹, Eldar Tavsultanovich Kaipaev², Daniil Rustamovich Vesenin^{3*}, Daniil Alexandrovich Tsarev⁴, Dmitry Egorovich Eremin⁴, Khamzat Saipudinovich Ikhaev⁵, Elvin Maarifovich Nuraliev⁶, Mata Ilyasovna Sakaeva¹, Anastasia Olegovna Popivnenko¹, Olga Alekseevna Karisheva¹, Olesya Andreevna Samofal⁶, Kirill Aleksandrovich Skrylnikov⁶

¹ North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, RUSSIA.

² Chechen State University named after A. A. Kadyrov, Grozny, Chechen Republic, RUSSIA.

³ Sechenov First Moscow State Medical University of the Ministry of Health of the Russian Federation (Sechenov University), Moscow, RUSSIA.

⁴ Ingush State University, Magas, Republic of Ingushetia, RUSSIA.

⁵ Kabardino-Balkarian State University named after H.M. Berbekov, Nalchik, Kabardino-Balkarian Republic, RUSSIA.

⁶ Rostov State Medical University, Rostov-on-Don, RUSSIA.

*Corresponding Author (Tel: +79183500889, ruslankalmykov777@yandex.ru).

Paper ID: 13A11P

Volume 13 Issue 11

Received 16 April 2022

Received in revised form 11

July 2022

Accepted 19 July 2022

Available online 25 July
2022

Keywords:

Anxiety; anxiety disorders; functional disorders; Gastrointestinal tract; Psychosomatic Aspects in Gastroenterology; Anxiety and depressive disorders; Depression cycle.

Abstract

Being widespread prevalence of anxiety disorders, one of the main causes of functional disorders of the gastrointestinal tract is the dysfunction of the nervous system. This review article discusses various cases of correction of functional disorders of the gastrointestinal tract. Based on many medical studies, 40% of patients with functional disorders of the gastrointestinal tract suffer from anxiety disorders. This study considers psychosomatic aspects in gastroenterology, as well as to study functional disorders of the gastrointestinal tract and their possible correction in patients with anxiety disorders. Due to the presence in many patients of gastroenterological profile, concomitant anxiety, psychopathic, psychovegetative, somatic disorders and manifestations, the use of psychopharmacotherapy is advisable and is often the only effective treatment option. The use of antidepressants, neuroleptics, tranquilizers, nootropics is carried out with strict consideration of the patient's condition and his individual characteristics.

Discipline: Medicine.

©2022 INT TRANS J ENG MANAG SCI TECH.

Cite This Article:

Makaeva, M. I., Khabalaeva, G. L., Kaipaev, E. T., Vesenin, D. R., Tsarev, D. A., Eremin, D. E., ... Skrylnikov K. A. (2022). Possibilities of Correction of Functional Diseases of the Gastrointestinal Tract in Patients with Anxiety Disorders. *International Transaction Journal of Engineering, Management, & Applied Sciences & Technologies*, 13(11), 13A11P, 1-10. <http://TUENGR.COM/V13/13A11P.pdf> DOI: 10.14456/ITJEMAST.2022.226

1 Introduction

Very often, patients have disorders of the gastrointestinal tract, but not all of them are accompanied by organic lesions, so most often such patients are diagnosed with functional gastrointestinal disorders (for example, functional dyspepsia, functional constipation or irritable bowel syndrome) [1-3].

The pathophysiology of functional disorders of the gastrointestinal tract is very complex. It includes a bidirectional violation of the regulation of the interaction of the intestine and the brain. The following conditions are attributed to the violation of the "intestine-brain" interaction: a change in the immune function of the intestinal mucosa, hypersensitivity, abnormal motility of the gastrointestinal tract, etc. [4].

Treatment of functional disorders of the gastrointestinal tract is not an easy task: underestimating the severity of psychoemotional disorders, such as anxiety or depression, leads to incorrect and, as a consequence, insufficiently effective treatment [5,6].

The principles of treatment are based on the study of the biopsychosocial foundations of the condition, including the therapy of physical symptoms and mental states. The purpose of the study is to consider psychosomatic aspects in gastroenterology, as well as to study functional disorders of the gastrointestinal tract and their possible correction in patients with anxiety disorders.

2 Psychosomatic Aspects in Gastroenterology

In recent decades, many scientists have been talking about an increase in the frequency of mental disorders. Anxiety disorders are most common among variants of borderline mental pathology. According to researchers, every fourth inhabitant of the planet had an anxiety disorder at least once in his life [7,8]. These disorders are found everywhere, and that is why this issue is very relevant.

Functional disorders of the gastrointestinal tract are more common in large cities and countries with developed economies [9]. One of the main causes of functional disorders of the gastrointestinal tract is dysfunction of the nervous system, frequent depression (Figure 1).




Figure 1. Depression cycle.

From an early age, a person is affected by many different factors: psychosocial, physiological and environmental factors, all of which often have a negative impact on the interaction of the central nervous system with the peripheral link of the nervous system, which is responsible for the work of the gastrointestinal tract. However, there are various causes of anxiety and depressive disorders (Table 1).

Table 1: Causes of anxiety and depressive disorders.

Type of disorders	Reasons	References
Psychogenic	They arise as a result of the impact on a person of an acute or chronic traumatic factor, including stress	[10,11]
Endogenous	The exact cause of the occurrence cannot be established, the presence of a genetic predisposition is likely	[12,13]
Somatogenic	Caused by the pathogenesis of another disease with somatic causes	[14,15]

The deterioration of the condition can contribute to: sedentary lifestyle, environmental degradation, multitasking, high mental and emotional stress, high pace of life, poor nutrition, these factors lead to functional disorders of the gastrointestinal tract. Thus, most patients are diagnosed with a psychosocial history that affects and disrupts the gastrointestinal tract-brain relationship [16].

Functional disorders that appear in a person at a young age progress over time and cause disorders in the psycho-emotional sphere, while further imbalance of the gastrointestinal tract is increasing. Often, with ineffective treatment, the patient's condition worsens, which is why, along with a functional disorder of the gastrointestinal tract, many concomitant diseases are detected in patients (for example, anxiety, depressive or hypochondriac disorders) [17-19]. The main symptoms of anxiety and depressive disorders are presented in Table 2

Table 2: The main symptoms of anxiety and depressive disorders

	Symptoms	Definition of the Symptom	References
Simple symptoms	Disturbing	Anxiety dominates, the appearance of phobias and panic attacks is possible	[20-23]
	Dreary	The main symptom of the disorder is a sharply lowered mood, a negative attitude to treatment, anhedonia	
	Dynamic	A general decrease in the dynamics of various processes, the appearance of motor and ideatory inhibition is characteristic	
	Anesthetic	Reduction of general physiological and emotional sensitivity, up to loss of sensitivity in certain parts of the body and the appearance of a feeling of painful insensibility	
Complex symptoms	With delirium	The appearance of various kinds of false beliefs that cannot be dissuaded, including hypochondriacal nature	[24-27]
	With hallucinations	The appearance of visual, auditory hallucinations, as well as senestopathies of various localization	
	Catatonic	Characterized by a significant decrease in mobility up to the state of stupor	
	Hidden	Hidden symptoms of anxiety and depressive symptoms	

Currently, the spread of the number of patients with functional gastrointestinal disorder is increasing, which is associated with an increase in the number of patients with psychosomatic pathology, which in turn is associated with impaired nervous system function [28,29].

Scientists and doctors associate the treatment of functional disorders of the gastrointestinal tract with disorders of nervous perception. Due to such conclusions, some definitions are recommended to be interpreted in a new way, for example, "functional disorder" is commonly called "violation of cerebro-intestinal interaction", and "functional abdominal pain" is now interpreted as "violation of the central perception of gastrointestinal pain" [30].

For the treatment of gastrointestinal motility, treatment with gastroenterological drugs is first recommended, with their ineffective effect, doctors recommend the use of antidepressants [8, 31, 32]. Currently, anti-anxiety drugs are more often used in practice, since they have a lower risk of side effects [33].

Thanks to numerous studies, it has become known that with functional disorders of the gastrointestinal tract, the activity of stress-limiting systems decreases, and the activity of the stress-releasing system begins to prevail. Because of this imbalance, anxiety states begin to form first, and then depression. Thus, it is possible to explain the symptoms that appear and develop with functional disorders of the gastrointestinal tract in the aggregate. These include: headaches, trembling, nervousness, irritability, sweating, dizziness, symptoms of dyspepsia and discomfort in the epigastric region [34-37].

3 Functional Disorders of the Gastrointestinal Tract and Their Possible Correction in Patients with Anxiety Disorders

A very common problem is insufficiently effective treatment of patients with functional disorders of the gastrointestinal tract (Table 1) [38,39]. The reason is an underestimation of the relationship between psychoemotional disorders, such as anxiety and depression, and the work of the gastrointestinal tract (Table 3).

Table 1. Factors influencing the development of functional disorders of the gastrointestinal tract.

Factors	Psychosocial aspects	Physiological aspects	References
Genetic	Stressful situations	Hormonal	[40-42]
Socio - cultural	Personality traits	Gastrointestinal sensitivity	[43-45]
Environmental	Psychological status	Immune dysfunction	[46-48]
	Susceptibility to stress	Microbiota change	
	Overcoming difficulties	Alimentary factor	
	Social support	Gastrointestinal motility	

For the prevention and treatment of psychosomatic disorders, doctors use complex treatment, including psychopharmacotherapy.

This therapy involves the solution of a number of tasks:

- identification and elimination of emotional disorders (anxiety, irritability, emotional instability);

- regulation of autonomic disorders;
- treatment and elimination of phobias;
- detection and leveling of negative factors;
- identification and elimination of somatic disorders.

In psychopharmacology, it is customary to distinguish the following groups of psychotropic drugs:

- neuroleptics;
- antidepressants;
- tranquilizers;
- nootropics.

From a large number of modern psychotropic drugs, it is necessary to choose those that have a wide range of efficacy (affect numerous symptoms); have a positive therapeutic effect; have a minimal set of side effects and insignificant severity of sedative effect [49,50].

Therefore, pharmacotherapy of psychosomatic diseases must meet the following requirements:

- be of a complex nature;
- combine psychotropic and somatotropic effects;
- take into account individual symptoms.

Thus, due to the presence in many patients of gastroenterological profile, concomitant anxiety, psychopathic, psychovegetative, somatic disorders and manifestations, the use of psychopharmacotherapy is advisable and is often the only effective treatment option. The use of antidepressants, neuroleptics, tranquilizers, nootropics is carried out with strict consideration of the patient's condition and his individual characteristics.

4 Conclusion

The gastrointestinal tract of a person has a close connection with his nervous system, therefore, the process of digestion and certain diseases of the gastrointestinal tract are affected by an imbalance of the sympathetic and parasympathetic departments. Disorders resulting from such an imbalance can have various symptoms (nausea, flatulence, diarrhea, constipation, etc.). Hence, there is a high level of coexistence of two diseases: functional disorders of the gastrointestinal tract and anxiety disorders.

Based on many medical studies, scientists conclude that about 40% of patients with functional disorders of the gastrointestinal tract suffer from anxiety disorders. The difficulty in diagnosing and distinguishing between somatoform and anxiety disorders is complicated due to the similarity of clinical manifestations. The former are bodily manifestations, and the latter are characterized by an emotional (mental) component.

That is why a gastroenterologist, when working with a patient with functional disorders of the gastrointestinal tract, should pay attention not only to the work of the digestive organs, but also look at the overall clinical picture of the disease. Only with such a versatile approach is the most correct and effective treatment of patients possible.

5 Availability of Data and Material

Data can be made available by contacting the corresponding author.

6 References

- [1] Beck A., Freeman A. Cognitive psychotherapy of personality disorders. St. Petersburg: Peter, 2019. 448 p.
- [2] Boraeva, T. T., Vadaeva, M. A., Matveeva, U. V., Revazova, A. B., Albegova, B. Z., Kanukoeva, D. T., Mishvelov, A. E. and Povetkin, S. N. Dynamics of Diseases of the Upper Digestive Tract in Children, *Journal of Pharmaceutical Research International*, 2021; 33(38B), 48-57. doi: 10.9734/jpri/2021/v33i38B32098
- [3] Blinov AV, Siddiqui SA, Nagdalian AA, Blinova AA, Gvozdenko AA, Raffa VV, et al. Investigation of the influence of Zinc-containing compounds on the components of the colloidal phase of milk. *Arab J Chem*. 2021;14(7):103229.
- [4] Drossman DA. Functional Gastrointestinal Disorders: History, Pathophysiology, Clinical Features and Rome IV. *Gastroenterology*. 2016:S0016-5085(16)00223-7. doi: 10.1053/j.gastro.2016.02.032
- [5] Lacy BE, Mearin F, Chang L, Chey WD, Lembo AJ, Simren M, Spiller, R. Bowel disorders. *Gastroenterology*. 2016;150:1393–1407.e5.
- [6] Khandia R. Codon Usage Analysis of Pro-Apoptotic Bim Gene Isoforms. 2022: 1711-1725
- [7] Maslova AY, Bazaeva KL, Abdullaeva ZA, Khazamova SO, Zeusheva KA, Grechkina TA, Semkina EN, Abramov MA, Mishvelov AE, Povetkin SN. Astrocytes and their Phenomenal Possibilities in the Treatment of Various Neurodegenerative Disorders: An Overview. *Journal of Pharmaceutical Research International*. 2021;33(33A): 60-68. DOI: 10.9734/jpri/2021/v33i33A31772
- [8] Siddiqui SA, Ali Redha A, Snoeck ER, Singh S, Simal-Gandara J, Ibrahim SA, Jafari SM. Anti-Depressant Properties of Crocin Molecules in Saffron. *Molecules*. 2022; 27(7):2076. DOI: 10.3390/molecules27072076
- [9] Firsova L.D. Generalized anxiety disorder in the practice of a gastroenterologist. Effective pharmacotherapy. 2020, 16(1): 80-83.
- [10] Donker, T., Griffiths, K.M., Cuijpers, P. *et al.* Psychoeducation for depression, anxiety and psychological distress: a meta-analysis. *BMC Med*. 2009;7,79. DOI: 10.1186/1741-7015-7-79
- [11] Bekhuis, Ella & Boschloo, Lynn & Rosmalen, Judith & Schoevers, Robert. Differential Associations of Specific Depressive and Anxiety Disorders with Somatic Symptoms. *Journal of Psychosomatic Research*. 2015; 78: 116-122. DOI: 10.1016/j.jpsychores.2014.11.007
- [12] Ozen D, Elif T. Symptom Similarities and Differences in Anxiety and Depressive Disorders. *Psikiyatride Guncel Yaklasimlar*. 2010; 2:1-15.
- [13] Goldberg D. Towards DSM-V: the relationship between generalized anxiety disorder and major depressive episode. *Psychol Med* 2008;38:1671-1675.
- [14] Fedosenko S. V., Kirillova N. A. The effect of treatment of severe chronic obstructive pulmonary disease with tiotropium bromide on comorbid somatogenic anxiety-depressive disorders. *Bulletin of Siberian Medicine*, 2011; 10(1):92-96.
- [15] Mykhaylov, V., Kozhyna, H., & Zdesenko, I. Psychotherapeutic correction system of somatogenic depressive disorders in patients with cerebral stroke. *INTER COLLEGAS*. 2017; 3(4): 190-196
- [16] Vyalov S.S. Factors of the pathogenesis of functional disorders and overlap syndrome of the gastrointestinal tract. *RMJ. Medical Review*. 2018; 7(2):68–71.
- [17] Musaeva PV, et al. Creation of New Types of Medical Simulation Systems with Feedback and

- [18] Bledzhyants GA, Mishvelov AE, Nuzhnaya KV, Anfinogenova OI, Isakova JA, Melkonyan RS. The Effectiveness of the medical decision-making support system “electronic clinical pharmacologist” in the management of patient’s therapeutic profile. *Pharmacophore.* 2019;10:76-81
- [19] Gvozdenko, A.A., Blinov, A.V., Slyadneva, K.S. *et al.* X-Ray Contrast Magnetic Diagnostic Tool Based on a Three-Component Nanosystem. *Russ J Gen Chem* 2022; 92:1153–1160. DOI: 10.1134/S1070363222060305
- [20] Craske, M., Stein, M., Eley, T. *et al.* Anxiety disorders. *Nat Rev Dis Primers.* 2017; 3, 17024. DOI: 10.1038/nrdp.2017.24
- [21] Health Quality Ontario. Internet-Delivered Cognitive Behavioural Therapy for Major Depression and Anxiety Disorders: A Health Technology Assessment. *Ont Health Technol Assess Ser.* 2019;19(6):1-199
- [22] Wingo TS, Liu Y, Gerasimov ES, Vattathil SM, Wynne ME, Liu J, Lori A, Faundez V, Bennett DA, Seyfried NT, Levey AI, Wingo AP. Shared mechanisms across the major psychiatric and neurodegenerative diseases. *Nat Commun.* 2022;13(1):4314. DOI: 10.1038/s41467-022-31873-5
- [23] Kouba BR, Camargo A, Gil-Mohapel J, Rodrigues ALS. Molecular Basis Underlying the Therapeutic Potential of Vitamin D for the Treatment of Depression and Anxiety. *Int J Mol Sci.* 2022 Jun 25;23(13):7077. DOI: 10.3390/ijms23137077
- [24] Millan MJ. Agomelatine for the treatment of generalized anxiety disorder: focus on its distinctive mechanism of action. *Ther Adv Psychopharmacol.* 2022;12:20451253221105128. doi: 10.1177/20451253221105128
- [25] Pérez de la Mora M, Borroto-Escuela DO, Crespo-Ramírez M, Rejón-Orantes JDC, Palacios-Lagunas DA, Martínez-Mata MK, Sánchez-Luna D, Tesoro-Cruz E, Fuxe K. Dysfunctional Heteroreceptor Complexes as Novel Targets for the Treatment of Major Depressive and Anxiety Disorders. *Cells.* 2022;11(11):1826. DOI: 10.3390/cells11111826
- [26] Al-Khazali HM, Ashina H, Iljazi A, Al-Sayegh Z, Lipton RB, Ashina M, Ashina S, Schytz HW. Psychiatric Sequelae Following Whiplash Injury: A Systematic Review. *Front Psychiatry.* 2022 Apr 22;13:814079. DOI: 10.3389/fpsy.2022.814079
- [27] Watanabe S, Omran AA, Shao AS, Xue C, Zhang Z, Zhang J, Davies DL, Shao XM, Watanabe J, Liang J. Dihydromyricetin improves social isolation-induced cognitive impairments and astrocytic changes in mice. *Sci Rep.* 2022 Apr 7;12(1):5899. DOI: 10.1038/s41598-022-09814-5
- [28] Vanner S, Greenwood-Van Meerveld B, Mawe G, Shea-Donohue T, Verdu EF, Wood J, Grundy D. Fundamentals of Neurogastroenterology: Basic Science. *Gastroenterology.* 2016 Feb 18:S0016-5085(16)00184-0. DOI: 10.1053/j.gastro.2016.02.018
- [29] Grinsvall C, Toˆrnblom H. Tack J Relationships between psychological state, abuse, somatization and visceral pain sensitivity in irritable bowel syndrome. *United European Gastroenterology Journal.* 2018; 6(2): 300–309.
- [30] Pichugina IM, Obraztsova VS, Frolov DI. Neurogastroenterology: benefits of interdisciplinary management of patients. *RMJ Medical Review.* 2018; 7(2):92–96.
- [31] Ayivi R, Ibrahim S, Colleran H, Silva R, Williams L, Galanakis C, Fidan H, Tomovska J and Siddiqui SA. COVID-19: human immune response and the influence of food ingredients and active compounds. *Bioactive Compounds in Health and Disease.* 2021;4(6):100

- [32] Profeta A, Siddiqui SA, Smetana S, Hossaini SM, Heinz V, Kircher C. The impact of Corona pandemic on consumer's food consumption: Vulnerability of households with children and income losses and change in sustainable consumption behavior. *J Verbrauch Lebensm.* 2021;16(4):305-314. DOI: 10.1007/s00003-021-01341-1
- [33] Rzhepakovsky IV, Areshidze DA, Avanesyan SS, Grimm WD, Filatova NV, Kalinin AV, Kochergin SG, Kozlova MA, Kurchenko VP, Sizonenko MN, Terentiev AA, Timchenko LD, Trigub MM, Nagdalian AA, Piskov SI. Phytochemical Characterization, Antioxidant Activity, and Cytotoxicity of Methanolic Leaf Extract of *Chlorophytum Comosum* (Green Type) (Thunb.) Jacq. *Molecules.* 2022; 27(3):762. DOI: 10.3390/molecules27030762
- [34] Raevskaya AI, Belyalova AA, Shevchenko PP, Karpov SM, Mishvelov AE, Simonov AN, Povetkin SN. et al. Cognitive Impairments in A Range of Somatic Diseases Diagnostics, Modern Approach to Therapy . *Pharmacophore.* 2020; 11(1):136-41.
- [35] Galabueva AI, Biragova AK, Kotsoyeva GA, Borukayeva ZK, Yesiev RK, Dzgoeva ZG, et al. Optimization of Modern Methods of Treating Chronic Generalized Periodontitis of Mild Severity. *Pharmacophore.* 2020;11(1):47-51.
- [36] Radovanovic-Dinica B., Tesic-Rajkovic S., Grgov S. Irritable bowel syndrome – from etiopathogenesis to therapy. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub.* 2018; 162(1):1-9.
- [37] Sobin WH, Heinrich TW, Drossman DA. Central Neuromodulators for Treating Functional GI Disorders: A Primer. *Am J Gastroenterol.* 2017;112(5):693-702. doi: 10.1038/ajg.2017.57
- [38] Van Oudenhove L, Crowell MD, Drossman DA, Halpert AD, Keefer L, Lackner JM, Murphy TB, Naliboff BD, Levy RL. Biopsychosocial Aspects of Functional Gastrointestinal Disorders. *Gastroenterology.* 2016 Feb 18:S0016-5085(16)00218-3. DOI: 10.1053/j.gastro.2016.02.027
- [39] Orsaeva AT, Tamrieva LA, Mischvelov AE, Osadchiy SS, Osipchuk GV, Povetkin SN, Simonov AN. Digital clinic “smart ward. *Pharmacophore.* 2020;11(1):142-146
- [40] Wu Y, Murray GK, Byrne EM. GWAS of peptic ulcer disease implicates *Helicobacter pylori* infection, other gastrointestinal disorders and depression. *Nat Commun.* 2021; 12:1146. DOI: 10.1038/s41467-021-21280-7
- [41] Akhmedov, Vadim. The development of functional gastrointestinal diseases: Genetic aspects. *Terapevticheskii arkhiv.* 2015; 87. 119-123. 10.17116/terarkh2015878119-123.
- [42] Villalba-Benito L, Torroglosa A, Luzón-Toro B, Fernández RM, Moya-Jiménez MJ, Antiñolo G, Borrego S. ChIP-Seq-Based Approach in Mouse Enteric Precursor Cells Reveals New Potential Genes with a Role in Enteric Nervous System Development and Hirschsprung Disease. *International journal of molecular sciences.* 2020;21(23):9061. DOI: 10.3390/ijms21239061
- [43] Holland AM, Bon-Frauches AC, Keszthelyi D, Melotte V, Boesmans W. The enteric nervous system in gastrointestinal disease etiology. *Cellular and Molecular Life Sciences.* 2021 May;78(10):4713-33. DOI: 10.1007/s00018-021-03812-y
- [44] *Ivashkin V.T., Shelygin Yu.A., Baranskaya E.K.* Clinical recommendations of the Russian Gastroenterological Association and the Association of Coloproctologists of Russia on the diagnosis and treatment of irritable bowel syndrome. *Ros journal gastroenterol hepatol coloproctol,* 2017; 27(5).
- [45] Esaulov V.I. Treatment of anxiety-phobic disorders in patients with irritable bowel syndrome by methods of integrative psychotherapy. *Modern problems of science and education.* 2017;1.

- [46] *Golovanova E.V.* Functional gastrointestinal disorders: approaches to the correction of psychosomatic disorders. *RMJ.* 2019; 27(5):24–29.
- [47] Baklanov IS, Baklanova OA, Shmatko AA, Gubanova MA, Pokhilko A.D. The historical past as a factor of sociocultural transformations of postmodernity. *Journal of History Culture and Art Research.* 2018; 7(1):373-378
- [48] Baklanov I, Rodionova V, Ivashova V, Shvachkina L, Medvedeva V. Social trends for increasing satisfaction with the educational services of a Modern University. *Quality-Access to Success.* 2020;21(179):88-90
- [49] Bottelier MA, Schouw ML, Klomp A. The effects of Psychotropic drugs On Developing brain (ePOD) study: methods and design. *BMC Psychiatry.* 2014; 14, 48. DOI: 10.1186/1471-244X-14-48
- [50] Ilyasov KK, Demchenkov EL, Chernyshkov AS, Rodin IA, Pushkin SV, Povetkin SN, et al. Features of the phytopharmacological preparations in the metaphylaxis of urolithiasis. *Pharmacophore.* 2020;11(5):66-71
-



Madina Georgievna Dzgoeva is a Student of North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Glikeria Levanovna Khabalaeva is a Student of North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Eldar Tavsultanovich Kaipaev is a Student of the Chechen State University named after A.A. Kadyrov, Grozny, Republic of Chechnya, Russia



Daniil Rustamovich Vesenin is a Student of the Moscow State Medical Sehenov, Moscow, Russia



Daniil Alexandrovich Tsarev is a Student of the General practitioner. Graduated from Yinggu Medical University.



Dmitry Egorovich Eremin is a Student of the Moscow State Medical Sehenov, Moscow, Russia



Elvin Maarifovich Nuraliev is a Student of Rostov State Medical University, Rostov-on-Don, Russia



Mata Ilyasovna Sakaeva is a Student of North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Khamzat Saipudinovich Ikhaev is a Student of Kabardino-Balkarian State University, Nalchik, Russia



Anastasia Olegovna Popivnenko is a Student of Rostov State Medical University, Rostov-on-Don, Russia



Olga Alekseevna Karisheva is a Student of North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Olesya Andreevna Samofal is a Student of Rostov State Medical University, Rostov-on-Don, Russia



Kirill Aleksandrovich Skrylnikov is a Student of Rostov State Medical University, Rostov-on-Don, Russia
