



The Role and Concept of Abdominoplasty at the Final Stages of Bariatric Interventions in Patients with Morbid Obesity

Anna Alexandrovna Yukhno¹, Ashura Gusengadzhievna Huseynova², Zaira Arazovna Abdullaeva², Zhamilat Arsenovna Abdullaeva², Deniza Aslanovna Vakalisheva³, Elena Alexandrovna Ulianova⁴, Alina Yurievna Maslova^{5,6}, Artem Evgenievich Mishvelov^{5,6}

¹ Kuban State Medical University, Krasnodar, RUSSIA

² Dagestan State Medical University, Makhachkala, Republic of Dagestan, RUSSIA

³ North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, RUSSIA

⁴ Tula State University, Tula, RUSSIA

⁵ Stavropol State Medical University, Stavropol, RUSSIA

⁶ Socmedica, Skolkovo, Moscow, RUSSIA

*Corresponding Author (Tel: +79183500889, Email: ruslankalmykov777@yandex.ru).

Paper ID: 13A8K

Volume 13 Issue 8

Received 26 January 2022

Received in revised form 28 May 2022

Accepted 04 June 2022

Available online 11 June 2022

Keywords:

Bariatric surgery; Morbid obesity; Body mass index; Abdominoplasty; Class III obesity; Massive weight loss; Post-bariatric treatment.

Abstract

Today we can say that morbid obesity is almost the main subject of such an urgent direction in medicine as bariatric surgery, which is indicated for patients with a BMI over 35 kg/m². This young rapidly developing area of surgical gastroenterology has not yet been sufficiently studied in the Russian Federation. However, according to the popularization of the method and the number of surgical interventions performed annually, this direction is rapidly gaining momentum. In comparison with 2003: the number of bariatric operations performed in Russia did not exceed 300-350 per year. According to the national bariatric registry for 2020, more than 16980 surgical interventions were performed in the Russian Federation. The National Bariatric Registry was created in 2013 on the initiative of the Stavropol doctor and President of the Society of Bariatric Surgeons of Russia concurrently - Bekhan Bayalovich Hatsiev, and it keeps records of absolutely all such operations and postoperative observations.

Disciplinary: Medicine.

©2022 INT TRANS J ENG MANAG SCI TECH.

Cite This Article:

Yukhno, A. A., Huseynova, A. G., Abdullaeva, Z. A., Abdullaeva, Zh. A., Vakalisheva, D. A., Ulianova, E. A., Maslova, A. Yu., Mishvelov, A. E. (2022) The Role and Concept of Abdominoplasty at the Final Stages of Bariatric Interventions in Patients with Morbid Obesity. *International Transaction Journal of Engineering, Management, & Applied Sciences & Technologies*, 13(8), 13A8K, 1-8. <http://TUENGR.COM/V13/13A8K.pdf> DOI: 10.14456/ITJEMAST.2022.158

1 Introduction

The most popular operation in Russia from 2013 to 2020 was longitudinal resection of the stomach (almost half of all types of operations). Gastric banding (the peak of popularity in 2013-2015) and mini gastric bypass surgery (actively gaining popularity recently) are located on the 2nd and 3rd places. 2020 was a record year for the number of entries in the national bariatric registry. In addition, this resource is one of the most complete among similar registries of other states. In 2020, he accumulated more than 90% of the total bariatric activity of the Russian Federation. This is the best indicator when compared with other national and international registries of bariatric surgery.

However, the numerous benefits that people receive after extremely rapid weight loss are accompanied by significant changes in appearance due to the formation of skin-fat folds throughout the body. Very often, such patients begin to turn to plastic surgeons, since otherwise, it is almost impossible to solve this problem [1, 2]. Yashkov, Yu.I. writes in his articles that "the situation should be considered optimal when a plastic surgeon works as part of a bariatric team" [3]. This review article discusses the role of abdominoplasty in the final stages of surgical treatment of patients with morbid obesity after bariatric surgery.

All this suggests that obesity surgery is proving its effectiveness. This review article discusses the role of abdominoplasty in the final stages of surgical treatment of patients with morbid obesity after bariatric surgery.

2 The Role of Abdominoplasty in the Treatment of Patients with Morbid Obesity

Over the past century and a half, the spectrum of diseases in the human community has been radically changed. If earlier people suffered and died mostly from infections, today there is a rampant epidemic of autoimmune disorders, allergies, insulin-dependent diabetes, and obesity. Moreover, the scale of the last two pathologies makes it necessary to take measures to eliminate them not only from the medical society but also at the state level in a number of countries. Although, as statistics and new data on the etiology and pathogenesis of obesity demonstrate, the problem is much more significant and deeper than it seemed before [1,4]. And in order to take certain steps towards correcting the situation in the context of modernity, there is a lot of research and social work to eliminate the causes and risk factors. The World Health Organization has officially dubbed obesity as a non-communicable epidemic that requires close attention, especially morbid obesity (hereinafter referred to as MO).

MO implies excessive deposition of fat mass, which is characterized by a BMI ≥ 40 kg/m² or a BMI ≥ 35 kg/m² in combination with serious complications associated with obesity [2]. That is, it can be a separate nosological unit and a motley syndrome with associated metabolic disorders [1,5]. This is one of the most serious causes of the development of arterial hypertension (hereinafter AH), sleep apnea syndrome, insulin-dependent diabetes, pathology of joints and spine, diseases of the digestive system, sexual disorders and even infertility. Obesity is accompanied by a decrease in

mental performance, causes psychological discomfort, a state of chronic depression and fatigue [2, 6].

These are well-known facts, but despite this, the scale of the problem is increasing. With the spread of MO, in parallel with the therapeutic and scientific side of solving the problem, a relatively young direction in surgery – bariatric surgery (hereinafter BS) began to develop actively. On the one hand, this happened due to the evidence of the ineffectiveness of conservative treatment in the long term in patients with MO: almost 95% of patients recover the lost mass within the first year [7-9], and insufficiently effective treatment of MO cannot be considered safe, since it does not eliminate the dangers of the disease itself. Therefore, we can safely say that MO is the main subject of BS, especially in economically developed countries, where, according to average estimates, MO occurs in 2-6% of the population [10].

The distribution of subcutaneous fat depends on many factors determined by gender and genetics. Male-type obesity involves the maximum accumulation of fat mass intraabdominal and under the skin of the anterior abdominal wall, in the pubic part. In women – in the hips, buttocks, mammary glands. The most common problem area is the anterior abdominal wall, where abdominoptoses are formed, known as "aprons". Patients with MO in the groin, on the anterior abdominal wall often develop a purulent rash, cellulite, and skin maceration, which are accompanied by wetness (wet rashes on the skin) and an unpleasant odor [11].

3 History of Abdominoplasty

The first abdominoplasty operation was performed in 1890 by Demarse and Marx, and the removal of a wide fat fold on the abdomen was developed by Kelly in 1899. At the beginning of the XX century, namely in 1939, Thorek described the complex abdominoplastic operations applicable to him in the work "Plastic removal of adipose tissue". Abdominoplasty is an operation that solves the problems of abdominoptosis, hanging of skin and fat folds throughout the body after extreme weight loss, postoperative deformed scars, flabby and weakened abdominal muscles, hernias, diastases, etc. In some sources, the term "abdominal dermatolipectomy" can be found, which means, as a rule, the elimination of an exclusively skin-fat flap without manipulations with the muscular-aponeurotic framework [2].

Due to the lack of awareness of the population about the possibilities of BS and the etiopathogenesis of their disease, many patients with MO turn to plastic surgeons to eliminate skin-fat "aprons", believing that this is an effective way to lose weight. In fact, abdominoplasty in such cases is a time-consuming and technically complex event due to the significant amount of wound discharge and the possibility of suppuration after surgery, and postoperative monitoring requires much more time and attention than the patient would turn to plastic surgery after bariatric surgery or self-weight loss. Ie, cosmetic results obtained against the background of uncorrected obesity cannot be considered either satisfactory or effective for the long term. In patients who have lost weight on their own or with the use of BS, abdominoplasty proceeds with less difficulties, both technical and restorative [11,12].

Types of operations possible with parallel BS operations:

1. Elimination of postoperative hernias;
2. Simultaneous elimination of other concomitant pathologies in the patient;
3. Simultaneous performance of the main bariatric intervention (according to an individual treatment plan) or simultaneous performance of corrective operations in other problem areas;
4. Elimination of complications and side effects of previous operations (for example, elongation of the small intestine in case of malabsorption syndrome) [2].

Accordingly, abdominoplasty cannot be used as a solution to MO and associated pathologies, but it can perfectly complement BS.

In patients with BS, abdominoplasty can be multi-purpose and multifunctional. In some cases, the elimination of excess skin-fat mass on the anterior abdominal wall is possible simultaneously with the main bariatric intervention [13]. Such an operation, panniculectomy, is used to increase the physical activity of patients by removing a large number of overhanging and often macerated flaps, i.e. it has a functional symptomatic character. At the same time, the surgical community warns that there is no need to expand the indications for such operations: it is necessary to be sure that abdominoplasty in a particular case can really seriously alleviate the patient's condition and benefit.

Another case where abdominoplasty can be applied, if the patient's condition allows, is a combination with restorative interventions on the small intestine. In rare cases, after gastric bypass surgery, there is a risk of developing protein deficiency and other nutritional deficiencies - due to excessive weight loss. If such a problem is not solved by conservative methods or it begins to be recurrent, then similar operations are used. Otherwise, when an insufficient result of the operation is noted against the background of weight stabilization, abdominoplasty can be combined with increased malabsorption due to shortening of the general and alimentary loops or with a further decrease in the volume of the stomach [14].

A complex of physical, laboratory and instrumental diagnostic studies make it possible to exclude symptomatic hypertension and identify possible lesions of other target organs [15-17]. The doctor should remember the need to use in practice the skills of objective research, including measurement and assessment of blood pressure, palpation of the pulse on the radial, carotid and femoral arteries, determination of the boundaries and tones of the heart, measurement of height and body weight (which, unfortunately, are often omitted, especially at the initial admission in the clinic) [18-22]. Such patients are advised to consult an optometrist to examine the fundus and detect retinal angioretinopathy. The list of mandatory studies includes a clinical blood test with platelet count, determination of the concentration of cholesterol, sugar, electrolytes, blood creatinine, general urinalysis, ECG registration. According to the indications, ultrasound examination of the kidneys, intravenous urography, renography with captopril (if renal artery stenosis is suspected), kidney angiography, echocardiography (EchoCG), and daily monitoring of blood pressure are performed. Along with an increase in blood pressure, the most important

diagnostic signs of primary arterial hypertension proper are a tense, firm pulse, expansion of the boundaries of the heart to the left due to an increase in the left ventricle, ECG and EchoCG signs of left ventricular hypertrophy, noise over the carotid arteries with their narrowing and in the epigastric region with renal artery stenosis, changes in the fundus (angioretinopathy retina) in the absence of symptoms of damage to the aorta and its large branches [23,24].

4 The Technique of Performing Abdominoplasty after Massive Weight Loss

On the day of the operation, the incisions and areas of the skin and subcutaneous tissue of the anterior abdominal wall are marked, which will be removed during the operation. Marking is better performed in a standing position since in the prone position the boundaries of the lateral excess flaps are lost. The lower incision line should provide correction of excess fat accumulations in the pubic area, and, as a rule, it is located 7, 0-10, 0 cm above the level of the genital slit in women.

Abdominoplasty is preferably performed under general anesthesia in the patient's back position. The peripheral vein is catheterized and a Foley catheter is installed. After the treatment of the surgical field, the skin is incised along the previously planned lines. Separately, the umbilical stalk is "cut out", which will be moved. The skin-fat flaps are mobilized 5-6 cm wider than the marking line for subsequent stitching of the flaps without tension. Then, after careful fitting and matching the edges of the incisions, excess skin-fat flaps are excised. To form the muscular-aponeurotic framework of the anterior abdominal wall, the aponeurosis is sutured along the midline with separate nodular sutures with non-absorbable threads. In the presence of postoperative hernias of the anterior abdominal wall, hernioplasty is performed more often using a mesh prosthesis, which is preferably located preperitoneal. Next, the navel is implanted into the newly marked place on the skin of the anterior abdominal wall. Subcutaneous sutures are applied with absorbable threads 3-0. Wound drainage is carried out by PCV tubes 5 mm in diameter through contrapertures in the suprapubic region. The operation is completed by applying an intradermal suture using absorbable threads.

At the first stages of the work, mainly transverse access or access in the form of an incomplete anchor is performed – dissection of the skin longitudinally upward to the navel level.

Advantages of Anchor access:

1. Creating an optimal contour of the abdomen with the formation of the waist;
2. Elimination of the massive "apron" creates a better visualization of the hernial defect and the edges of aponeurosis, which facilitates hernioplasty if necessary;
3. Free access to any part of the abdominal cavity through a hernial defect of the anterior abdominal wall or through a separate mini-laparotomy;

Wide excision of excess skin-fat tissue provides free access to any part of the abdominal cavity. If there was a hernia, then access to the abdominal organs was carried out through a hernial

defect of the anterior abdominal wall. From such access, it is quite convenient to perform longitudinal resection of the stomach, as well as any operation on the small intestine.

In practice, surgeons try to separate the edges of the flaps as little as possible in the area of the median incision and in the area of the lateral flaps. This helps to preserve the blood supply to the lateral flaps, preventing the development of marginal necrosis. This prevents the development of gray postoperative wounds. In order to strengthen the weak points of the anterior abdominal wall, the duplication of aponeurosis is also formed by monofilament threads with long resorption periods [12-15].

5 Conclusion

Bariatric Surgery (BS) is a young and rapidly developing area in obesity surgery in Russia. Abdominoplasty surgery can play a useful role in improving cosmetic results and the functional component of the life of patients with MO in combination with bariatric operations. However, it should be remembered that cosmetic results obtained against the background of uncorrected obesity cannot be considered either satisfactory or effective in the long term. In patients who have lost weight on their own or with the use of BS, abdominoplasty proceeds with less difficulties, both technical and restorative. The treatment of "post-bariatric" patients is a difficult task for a doctor, since it requires a step-by-step approach to choosing tactics, performing complex accesses, and using various high-tech surgical techniques. Careful planning of the operation allows you to minimize side effects and complications.

6 Availability of Data and Material

Data can be made available by contacting the corresponding authors.

7 References

- [1] Collen A, Hanemaaijer R, Lupu F, Quax PH, van Lent N, Grimbergen J, Peters E, Koolwijk P, van Hinsbergh VW. Membrane-type matrix metalloproteinase-mediated angiogenesis in a fibrin-collagen matrix. *Blood*. 2003 Mar 1;101(5):1810-7. doi: 10.1182/blood-2002-05-1593.
- [2] Richards TA, Kaye AD, Fields AM. Morbid obesity--a review. *Middle East J Anaesthesiol*. 2005 Feb;18(1):93-105.
- [3] Dedov II, Melnichenko GA, Troshina EA, Ershova EV, Mazurina NV, Ogneva NA, Yashkov YI, Ilyin AV. Produktsiya inkretinov i obmen glyukozy u bol'nykh patologicheskim ozhireniem v rannem i otdalennom periode posle biliopankreaticheskogo shuntirovaniya [Incretin secretion and glucose metabolism in morbidly obese patients in the early and late periods after biliopancreatic diversion]. *Ter Arkh*. 2016;88(10):9-18. Russian. doi: 10.17116/terarkh201688109-18
- [4] Siddiqui Shahida Anusha, Redha Ali Ali, Esmaeili Yasaman & Mehdizadeh Mohammad. Novel insights on extraction and encapsulation techniques of elderberry bioactive compounds, *Critical Reviews in Food Science and Nutrition*, 2002, DOI: 10.1080/10408398.2022.2026290
- [5] Ranjha MMAN, Shafique B, Rehman A, Mehmood A, Ali A, Zahra SM, Roobab U, Singh A, Ibrahim SA and Siddiqui SA (2022) Biocompatible Nanomaterials in Food Science, Technology, and Nutrient Drug Delivery: Recent Developments and Applications. *Front. Nutr.* 8:778155. doi: 10.3389/fnut.2021.778155
- [6] Shahida Anusha Siddiqui, Nur Alim Bahmid, Ahmed Taha, Ibrahim Khalifa, Sipper Khan, Hadis

Rostamabadi & Seid Mahdi Jafari (2022) Recent advances in food applications of phenolic-loaded micro/nanodelivery systems, *Critical Reviews in Food Science and Nutrition*, DOI: 10.1080/10408398.2022.2056870

- [7] Benaiges D, Goday A, Pedro-Botet J, Más A, Chillarón JJ, Flores-Le Roux JA. Bariatric surgery: to whom and when? *Minerva Endocrinol.* 2015 Jun;40(2):119-28.
- [8] Bozola AR, Psillakis JM. Abdominoplasty: a new concept and classification for treatment. *Plast Reconstr Surg.* 1988 Dec;82(6):983-93.
- [9] Christman KD. Death following suction lipectomy and abdominoplasty. *Plast Reconstr Surg.* 1986 Sep;78(3):428. doi: 10.1097/00006534-198609000-00040.
- [10] Dubou R, Ousterhout DK. Placement of the umbilicus in an abdominoplasty. *Plast Reconstr Surg.* 1978 Feb;61(2):291-3. doi: 10.1097/00006534-197802000-00030.
- [11] Flageul G, Elbaz JS, Karcenty B. Les complications de la chirurgie plastique de l'abdomen [Complications of plastic surgery of the abdomen]. *Ann Chir Plast Esthet.* 1999 Aug;44(4):497-505. French.
- [12] Cabbabe SW. Plastic Surgery after Massive Weight Loss. *Mo Med.* 2016 May-Jun;113(3):202-6.
- [13] Floros C, Davis PK. Complications and long-term results following abdominoplasty: a retrospective study. *Br J Plast Surg.* 1991 Apr;44(3):190-4. doi: 10.1016/0007-1226(91)90125-4
- [14] Hakme F. Abdominoplasty: peri- and supra-umbilical lipectomy. *Aesthetic Plast Surg.* 1983;7(4):213-20. doi: 10.1007/BF01570662
- [15] Tovlahanova TJH, et al. Study of the Effect of the Image Scanning Speed and the Type of Conductive Coating on the Quality of Sem-Micrographs of Oxide Nano Materials for Medical Use. *Ann Med Health Sci Res.* 2021;11,S3:60-64
- [16] Rzhepakovsky I, Anusha Siddiqui S, Avanesyan S, Benlidayi M, Dhingra K, Dolgalev A, Erukashvily N, Fritsch T, Heinz V, Kochergin S, Nagdalian A, Sizonenko M, Timchenko L, Vukovic M, Piskov S, Grimm WD. Anti-arthritic effect of chicken embryo tissue hydrolyzate against adjuvant arthritis in rats (X-ray microtomographic and histopathological analysis). *Food Sci Nutr.* 2021 Aug 18;9(10):5648-5669. doi: 10.1002/fsn3.2529
- [17] Bazhenova AA, Guryanova NI, Guryanov GS, Alieva HAV, Kachmazova DT, Khripunova A A, et al. In-Vitro Study of the Properties of Components for the Synthesis of Sorbent for Low-Density Lipoprotein Apheresis. *Pharmacophore.* 2021;12(3):37-41. DOI: 10.51847/BsjhKFW0Kd
- [18] Ramirez OM. Abdominoplasty and abdominal wall rehabilitation: a comprehensive approach. *Plast Reconstr Surg.* 2000 Jan;105(1):425-35. doi: 10.1097/00006534-200001000-00069
- [19] Hight G. Ya, Mishvelov AE, Nuzhnaya CV et al. New image modeling features for planning surgical interventions. *Research Journal of Pharmaceutical, Biological and Chemical Sciences.* 2019;10(1):140-143
- [20] Hite GJ, Mishvelov AE, Melchenko EA, Vlasov AA, Anfinogenova OI, Nuzhnaya CV, et al. Holodoctor planning software real-time surgical intervention. *Pharmacophore.* 2019;10(3):57-60
- [21] Bouyahya A, El Menyiy N, Oumeslakht L, El Allam A, Balahbib A, Rauf A, Muhammad N, Kuznetsova E, Derkho M, Thiruvengadam M, Shariati MA, El Omari N. Preclinical and Clinical Antioxidant Effects of Natural Compounds against Oxidative Stress-Induced Epigenetic Instability in Tumor Cells. *Antioxidants.* 2021; 10(10):1553. <https://doi.org/10.3390/antiox10101553>
- [22] Ya S. Shevchenko, Plohova DP, Bulakhova IN, Mishvelov AE, Kubalova ME, Badriev GB, Kh. A.

Mildzikhov, Simonov AN, Verevkina MN, Okolelova AI, Povetkin SN. Experience of carrying out magnetic resonance imaging with the use of specialized protocols and programs computer post-processing. Pharmacophore. 2020;11(2):77-81

[23] Teimourian B. Tourniquet after suction lipectomy of the lower extremity. Plast Reconstr Surg. 1985 Mar;75(3):442. doi: 10.1097/00006534-198503000-00040.

[24] Matarasso A, Aly A, Hurwitz DJ, Lockwood TE. Body contouring after massive weight loss. Aesthet Surg J. 2004 Sep-Oct;24(5):452-63. doi: 10.1016/j.asj.2004.04.006.




Anna Alexandrovna Yukhno is a student of Kuban State Medical University, Krasnodar, Russia.




Ashura Gusengadzhievna Huseynova is a student of Dagestan State Medical University, Makhachkala, Republic of Dagestan, Russia.




Zaira Arazovna Abdullaeva is a student of Dagestan State Medical University, Makhachkala, Republic of Dagestan, Russia.



Zhamilat Arsenovna Abdullaeva is a student of Dagestan State Medical University, Makhachkala, Republic of Dagestan, Russia



Deniza Aslanovna Vakalisheva is a student of North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Elena Alexandrovna Ulianova is a student of Tula State University, Tula, Russia.



Dr. Alina Yurievna Maslova is a Junior Researcher at the Pharmacological Laboratory of Stavropol State Medical University, Stavropol, Russia. Medical Expert, Socmedica, Skolkovo, Moscow, Russia.



Dr. Artem Evgenievich Mishvelov is a Junior Researcher at the 3D Technology Laboratory of Stavropol State Medical University, Stavropol, Russia. Head of Project Activities in the North Caucasus Federal District, Socmedica, Skolkovo, Moscow, Russia
