



The Effectiveness of the Diagnosis of Endothelial Dysfunction and its Pharmacological Correction with a Focus on Angiotensin-Converting Enzyme Inhibitors

Ilona Viktorovna Bestaeva¹, Murad Aslambekovich Khasanov², Gulshan Anver Babirova³, Zakhra Zauravna Askerova³, Lora Khasanbekovna Dzotsieva¹, Alina Yurievna Maslova^{4,5}

¹ North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, RUSSIA.

² Chechen State University, Grozny, Republic of Chechnya, RUSSIA.

³ Dagestan State Medical University, Makhachkala, Republic of Dagestan, RUSSIA.

⁴ Stavropol State Medical University, Stavropol, RUSSIA.

⁵ Socmedica, Skolkovo, Moscow, RUSSIA.

*Corresponding Author (Tel: +79183500889, Email: ruslankalmykov777@yandex.ru).

Paper ID: 13A8L

Volume 13 Issue 8

Received 21 January 2022

Received in revised form 31

May 2022

Accepted 07 June 2022

Available online 14 June

2022

Keywords:

Arterial hypertension;
Endothelial dysfunction;
Angiotensin-converting
enzyme inhibitors;
Vascular endothelium;
Nitrogen monoxide;
Oxidative stress.

Abstract

The vascular endothelial lining is a kind of unique endocrine system, which is located around the perimeter of absolutely all organs of the vascular system. Why endocrine: in modern cardiology, the main importance in the occurrence and development of almost all pathologies is assigned to neurohormonal breakdowns in the form of an imbalance between vasoconstrictors and vasodilators. On one side of the hormonal balance are neurohormones that cause vasoconstriction, remodeling and antidiuresis. These are components of the renin-angiotensin-aldosterone and sympathoadrenal systems, endothelin, vasopressin. They are counteracted by hormones with vasodilating and diuretic effects that block remodeling processes, the main of which are nitrogen monoxide, as well as natriuretic peptides, the kallikrein-kinin system and prostacyclin. Endothelial cells play the role of an effective buffer between the blood flow and the tissues of the body, performing a number of regulatory functions, producing a large variety of biologically active substances. The strategic location of the endothelium allows it to be sensitive to the slightest changes in the hemodynamic system and not only in it. Thus, the vascular endothelium plays a key role in maintaining homeostasis. This review article presents the main pathogenetic aspects of endothelial dysfunction, methods of diagnosis and its correction.

Disciplinary: Medicine.

©2022 INT TRANS J ENG MANAG SCI TECH.

Cite This Article:

Bestaeva, I.V., Khasanov, M. A., Babirova, G. A., Askerova, Z. Z., Dzotsieva, L. K., Maslova, A.Yu. (2022). The effectiveness of the diagnosis of endothelial dysfunction and its pharmacological correction with a focus on angiotensin-converting enzyme inhibitors. *International Transaction Journal of Engineering, Management, & Applied Sciences & Technologies*, 13(8), 13A8L, 1-8. <http://TUENGR.COM/V13/13A8L.pdf> DOI: 10.14456/ITJEMAST.2022.159

1 Introduction

In classical histology and physiology, the vascular endothelium layer is a layer of flat cells of mesenchymal origin lining the inner surface of blood and lymphatic vessels [1]. The endothelium is not just a semipermeable membrane, but an active endocrine organ. After discovering and studying the functions of nitrogen monoxide as a signaling molecule of the vascular system, scientists were able to pave a new direction in understanding the involvement of the endothelium in the pathogenesis of hypertension and other cardiovascular pathologies, as well as ways to effectively correct its dysfunction [2].

The most important functions of the endothelium are the maintenance of hemovascular homeostasis, regulation of hemostasis, modulation of inflammation, regulation of vascular tone and vascular permeability. In addition, the endothelium has its own renin-angiotensin system. The endothelium secretes mitogens and participates in angiogenesis, fluid balance, and the exchange of intercellular matrix components [3-5]. The vascular endothelium performs these functions by synthesizing and isolating a large number of various biologically active substances (Table 1) [6-10].

Table 1: Factors synthesized in the endothelium and determining its functions

Factors affecting vascular smooth muscle tone	
Vasoconstrictors	Vasodilators
Endothelin	Nitrogen monoxide
Angiotensin II	Prostacyclin
Thromboxane A2	Endothelin depolarization factor
Prostaglandin H2	Angiotensin I Adrenomedullin
Hemostasis factors	
Prothrombogenic	Antithrombogenic
Platelet growth factor	Thrombomodulin
Inhibitor of tissue plasminogen activator	Nitrogen monoxide
Angiotensin IV	Tissue Plasminogen activator
Platelet activation factor	Prostacyclin
The Willebrand Factor	
Endothelin I	
Fibronectin	
Thrombospondin	
Factors affecting growth and proliferation	
Stimulants	Inhibitors
Endothelin I	Nitrogen monoxide
Angiotensin II	Prostacyclin
Superoxide radicals	Natriuretic C-type peptide
Endothelial Growth Factor	Heparin-like growth inhibitors
Factors affecting inflammation	
Pro - inflammatory	Anti-inflammatory
Tumor necrosis factor alpha	Nitrogen monoxide
Superoxide radicals	
C-reactive protein	

2 Pathophysiological Aspects of the Development of Endothelial Dysfunction

The main task of the endothelium is the balanced release of biologically active substances that determine the holistic work of the circulatory system [11]. In the physiological state, the

endothelium has the ability to maintain a balance between its multidirectional functions: synthesis of pro- and anti-inflammatory factors, vasodilating and vasoconstrictive substances, pro- and antiplatelet agents, pro- and anticoagulants, pro- and antifibrinolytics, proliferation factors and growth inhibitors [12-14]. Under physiological conditions, vasodilation, synthesis of aggregation inhibitors, coagulation and fibrinolysis activators, anti-adhesive substances prevail. Vascular cell dysfunction disrupts this balance and predisposes vessels to vasoconstriction, leukocyte adhesion, platelet activation, mitogenesis and inflammation [13]. Thus, the endothelial function is a balance of oppositely acting principles: relaxing and constrictive factors, anticoagulant and procoagulant factors, growth factors, and their inhibitors [14].

Causes such as impaired blood flow, hypoxia, increased systemic and intrarenal pressure, hyperhomocysteinemia, and increased lipid peroxidation processes can lead to changes in the physiological balance in the body [15-17]. Vascular endothelium is extremely vulnerable, but, on the other hand, researchers note its huge compensatory capabilities in violation of physiological conditions [18].

Currently, endothelial dysfunction is understood as an imbalance between the formation of vasodilating, atrombogenic, antiproliferative factors, on the one hand, and vasoconstrictive, prothrombotic and proliferative substances synthesized by the endothelium, on the other. Endothelial dysfunction can be an independent cause of circulatory disorders in the organ, since it often provokes angiospasm or vascular thrombosis. On the other hand, disorders of regional blood circulation (ischemia, venous congestion) can also lead to endothelial dysfunction [19]. Hemodynamic causes, age-related changes, free radical damage, dyslipoproteinemia, hypercytokinemia, hyperhomocysteinemia, exogenous and endogenous intoxication can contribute to the formation of endothelial dysfunction [20]. Endothelial dysfunction can lead to structural damage in the body: acceleration of apoptosis, necrosis, desquamation of endotheliocytes. However, functional changes in the endothelium, as a rule, precede morphological changes in the vascular wall [21].

There are four forms of endothelial dysfunction: vasomotor, thrombophilic, adhesive and angiogenic [22]. The vasomotor form of endothelial dysfunction is caused by a violation of the relationship between endothelial vasoconstrictors and vasodilators and is important in the mechanisms of both systemic increases in blood pressure and local angiospasm. The thrombophilic form of endothelial dysfunction is caused by a violation of the ratio of thrombogenic and atrombogenic substances formed in the endothelium and involved in hemostasis or affecting this process. Under physiological conditions, the formation of atrombogenic substances in the endothelium prevails over the formation of thrombogenic substances, which ensures the preservation of the liquid state of the blood in case of damage to the vascular wall. The thrombophilic form of endothelial dysfunction can lead to the development of vascular thrombophilia and thrombosis. The adhesive form of endothelial dysfunction is caused by a violation of the interaction of leukocytes and endothelium. Increased adhesion of the endothelium

and uncontrolled adhesion of leukocytes are of great importance in the pathogenesis of inflammation in atherosclerosis and other pathological processes. The angiogenic form of endothelial dysfunction is associated with a violation of neoangiogenesis. Therefore, endothelial dysfunction as a typical pathological process is a key link in the pathogenesis of many diseases and their complications [23,24].

3 Diagnosis of Endothelial Dysfunction

Direct determination of NO in the blood is difficult due to its short half-life. NO production is evaluated by the content in blood plasma and urine of the end products of NO metabolism (nitrite, nitrate) during chemiluminescence analysis or gas chromatography/mass spectrometry [25,26]. The determination of cyclic guanosine monophosphate (cGMP) makes it possible to estimate the proportion of NO that had a functional effect on endothelial cells and to distinguish reduced NO production from increased degradation. The concentrations of these substances are extremely low, there is a significant dependence on other sources of nitrite and nitrate, including food, and therefore the clinical use of this method is limited. Endothelin-1 is an endothelial peptide with strong vasoconstrictor, mitogenic properties. It is believed that an increased level of endothelin-1 is associated with endotheliocyte damage and can be considered a marker of ED [27]. Due to large fluctuations in people, regardless of vascular status, the use of endothelin-1 as a separate marker is highly inaccurate. Willebrand factor is a glycoprotein synthesized mainly by endothelial cells. Its level correlates with the severity of risk factors for atherosclerosis, the elimination of which is accompanied by its decrease. An increased level of the Willebrand factor indicates either endothelial damage or, more likely, increased activation of endotheliocytes [28]. Tissue plasminogen activator, plasminogen activator inhibitor-1: with endothelial dysfunction and an increased risk of atherosclerosis or its complications, an increase in the level of both of these markers is observed. Adhesive molecules include vascular cellular adhesive molecule 1 (VCAM 1), endothelial-leukocyte adhesive molecule 1 (E-selectin), inter-cellular adhesive molecule 1 (ICAM 1), and P-selectin. Circulating soluble forms of adhesive molecules are detected in plasma, their levels are elevated in inflammatory diseases, and they are measured by immunological methods.

4 Pharmacological Correction of Endothelial Dysfunction Based on Angiotensin Converting Enzyme (ACE) Inhibitors

At least two main mechanisms of ACEI influence on endothelial function are discussed. It is known that ACE inhibitors lead to an increase in the amount of tissue bradykinin. Vasoprotective, antiproliferative, antisclerotic and acute vasodilatory effects of ACE inhibitors can be explained by endothelium-dependent reactions associated with the ability of ACE inhibitors to prevent the breakdown of bradykinin. Bradykinin is a powerful stimulant for the release of endothelium-dependent relaxing factors such as NO, endothelium-dependent hyperpolarization factor and prostacyclin. Endothelial hyperpolarization factor is an unstable metabolite of arachidonic acid, its release depends on the concentration of intracellular calcium and calmodulin, and its effect on endothelium-dependent relaxation is associated with the size of blood vessels and is most

significant in small arteries. Prostacyclin, formed in endothelial cells, activates adenylate cyclase of smooth muscle cells, increasing the formation of cAMP, the relaxing effect of which enhances vasodilation caused by NO. The concept of the possibility of using ACE inhibitors to affect endothelial function, ischemia and atherogenesis has emerged relatively recently. This effect is realized with the help of several mechanisms. The first information that ACE inhibitors contribute to a decrease in the frequency of myocardial ischemic events was obtained as a result of large clinical studies involving patients with left ventricular dysfunction and heart failure. At the end of these studies, it became obvious that the use of ACE inhibitors contributes to a significant reduction in the number of ischemic negative events [29].

Another mechanism of action of ACE inhibitors on endothelial function is the blockade of angiotensin II formation, which is considered an inducer of oxidative stress. An increase in the formation of reactive oxygen species (superoxide anion) has been experimentally demonstrated under the action of angiotensin II. The mechanisms of action of angiotensin II on the production of superoxide anion are associated with the stimulation of NADPH/NADH oxidases [30,31]. Thus, a decrease in the level of angiotensin II leads to a decrease in oxidative stress, the products of which reduce the activity of NO. Angiotensin II in many respects has the opposite effect with respect to NO and is currently recognized as practically its antagonist. ACE inhibition restores the balance between the two vasoactive systems [32].

5 Conclusion

To date, the concept of endothelial dysfunction as a central link in the pathogenesis of many chronic diseases has been formulated. The main role in the development of endothelial dysfunction is played by oxidative stress, the synthesis of powerful vasoconstrictors that suppress the formation of nitrogen monoxide. Endothelial dysfunction precedes the development of clinical manifestations of diseases, therefore, the evaluation of endothelial functions is of great diagnostic and prognostic importance. Further study of the role of endothelial dysfunction in the development of diseases is necessary to develop new therapeutic approaches. Currently, there is an accumulation of new data on the specific mechanisms of endothelial dysfunction in hypertension. The effect of traditional antihypertensive drugs is supplemented and reinterpreted from the point of view of the effect on the endothelium, and the mechanisms of vasoprotective and organoprotective action of cardiovascular drugs due to the modulation of NO production are revealed. Priority endothelium-protective drugs today are ACE inhibitors with the maximum level of lipophilicity.

6 Availability of Data and Material

Data can be made available by contacting the corresponding authors.

7 References

- [1] Konukoglu D, Uzun H. Endothelial Dysfunction and Hypertension. *Adv Exp Med Biol.* 2017;956:511-540. DOI: 10.1007/5584_2016_90.
- [2] Tovlahanova TJH, et al. Study of the Effect of the Image Scanning Speed and the Type of Conductive Coating on the Quality of Sem-Micrographs of Oxide Nano Materials for Medical Use. *Ann Med Health*

- [3] Annuk M, Zilmer M, Lind L, Linde T, Fellström B. Oxidative stress and endothelial function in chronic renal failure. *J Am Soc Nephrol.* 2001 Dec;12(12):2747-2752. DOI: 10.1681/ASN.V12122747.
- [4] Lee JM, Layland J, Jung JH, Lee HJ, Echavarría-Pinto M, Watkins S, Yong AS, Doh JH, Nam CW, Shin ES, Koo BK, Ng MK, Escaned J, Fearon WF, Oldroyd KG. Integrated physiologic assessment of ischemic heart disease in real-world practice using index of microcirculatory resistance and fractional flow reserve: insights from the International Index of Microcirculatory Resistance Registry. *Circ Cardiovasc Interv.* 2015 Nov;8(11):e002857. DOI: 10.1161/CIRCINTERVENTIONS.115.002857.
- [5] Guzik TJ, Harrison DG. Vascular NADPH oxidases as drug targets for novel antioxidant strategies. *Drug Discov Today.* 2006 Jun;11(11-12):524-33. DOI: 10.1016/j.drudis.2006.04.003.
- [6] Noma K, Goto C, Nishioka K, Jitsuiki D, Umemura T, Ueda K, Kimura M, Nakagawa K, Oshima T, Chayama K, Yoshizumi M, Liao JK, Higashi Y. Roles of rho-associated kinase and oxidative stress in the pathogenesis of aortic stiffness. *J Am Coll Cardiol.* 2007 Feb 13;49(6):698-705. DOI: 10.1016/j.jacc.2006.06.082.
- [7] Naceiri Mrabti, Hanae & Bouyahya, Abdelhakim & Ed-Dra, Abdelaziz & Kachmar, Mohamed Reda & Naceiri Mrabti, Nidal & Benali, Taoufiq & Shariati, Mohammad Ali & Ouahbi, Abdelilah & Doudach, Latifa & my el abbes, Faouzi. (2021). Polyphenolic profile and biological properties of *Arbutus unedo* root extracts. *European Journal of Integrative Medicine.* 42. 10.1016/j.eujim.2020.101266
- [8] Siddiqui Shahida Anusha, Redha Ali Ali, Esmaeili Yasaman & Mehdizadeh Mohammad. Novel insights on extraction and encapsulation techniques of elderberry bioactive compounds, *Critical Reviews in Food Science and Nutrition*, 2022, DOI: 10.1080/10408398.2022.2026290
- [9] Bouyahya A, El Hachlafi N, Aanniz T, Bourais I, Mechchate H, Benali T, Shariati MA, Burkov P, Lorenzo JM, Wilairatana P, Mubarak MS, El Omari N. Natural Bioactive Compounds Targeting Histone Deacetylases in Human Cancers: Recent Updates. *Molecules.* 2022; 27(8):2568. <https://doi.org/10.3390/molecules27082568>
- [10] Mechchate H, El Allam A, El Omari N, El Hachlafi N, Shariati MA, Wilairatana P, Mubarak MS, Bouyahya A. Vegetables and Their Bioactive Compounds as Anti-Aging Drugs. *Molecules.* 2022; 27(7):2316. <https://doi.org/10.3390/molecules27072316>
- [11] Thiruvengadam M, Venkidasamy B, Subramanian U, Samynathan R, Ali Shariati M, Rebezov M, Girish S, Thangavel S, Dhanapal AR, Fedoseeva N, Lee J, Chung I-M. Bioactive Compounds in Oxidative Stress-Mediated Diseases: Targeting the NRF2/ARE Signaling Pathway and Epigenetic Regulation. *Antioxidants.* 2021; 10(12):1859. <https://doi.org/10.3390/antiox10121859>
- [12] Shahida Anusha Siddiqui, Nur Alim Bahmid, Ahmed Taha, Ibrahim Khalifa, Sipper Khan, Hadis Rostamabadi & Seid Mahdi Jafari (2022) Recent advances in food applications of phenolic-loaded micro/nanodelivery systems, *Critical Reviews in Food Science and Nutrition*, DOI: 10.1080/10408398.2022.2056870
- [13] Ranjha MMAN, Shafique B, Rehman A, Mehmood A, Ali A, Zahra SM, Roobab U, Singh A, Ibrahim SA and Siddiqui SA (2022) Biocompatible Nanomaterials in Food Science, Technology, and Nutrient Drug Delivery: Recent Developments and Applications. *Front. Nutr.* 8:778155. DOI: 10.3389/fnut.2021.778155
- [14] Rzhepakovsky IV, Areshidze DA, Avanesyan SS, Grimm WD, Filatova NV, Kalinin AV, Kochergin SG, Kozlova MA, Kurchenko VP, Sizonenko MN, Terentiev AA, Timchenko LD, Trigub MM, Nagdalian AA, Piskov SI. Phytochemical Characterization, Antioxidant Activity, and Cytotoxicity of Methanolic Leaf Extract of *Chlorophytum Comosum* (Green Type) (Thunb.) Jacq. *Molecules.* 2022; 27(3):762.


- [15] Querfeld U, Mak RH, Pries AR. Microvascular disease in chronic kidney disease: the base of the iceberg in cardiovascular comorbidity. *Clin Sci (Lond)*. 2020 Jun 26;134(12):1333-1356. DOI: 10.1042/CS20200279.
- [16] Rzhepakovsky I, Anusha Siddiqui S, Avanesyan S, Benlidayi M, Dhingra K, Dolgalev A, Enuakashvily N, Fritsch T, Heinz V, Kochergin S, Nagdalian A, Sizonenko M, Timchenko L, Vukovic M, Piskov S, Grimm WD. Anti-arthritic effect of chicken embryo tissue hydrolyzate against adjuvant arthritis in rats (X-ray microtomographic and histopathological analysis). *Food Sci Nutr*. 2021 Aug 18;9(10):5648-5669. DOI: 10.1002/fsn3.2529
- [17] Bouyahya A, El Menyiy N, Oumeslakht L, El Allam A, Balahbib A, Rauf A, Muhammad N, Kuznetsova E, Derkho M, Thiruvengadam M, Shariati MA, El Omari N. Preclinical and Clinical Antioxidant Effects of Natural Compounds against Oxidative Stress-Induced Epigenetic Instability in Tumor Cells. *Antioxidants*. 2021; 10(10):1553. DOI: 10.3390/antiox10101553
- [18] Deanfield JE, Halcox JP, Rabelink TJ. Endothelial function and dysfunction: testing and clinical relevance. *Circulation*. 2007 Mar 13;115(10):1285-95. DOI: 10.1161/CIRCULATIONAHA.106.652859
- [19] Kuryliszyn-Moskal A, Klimiuk PA, Sierakowski S, Ciolkiewicz M. A study on vascular endothelial growth factor and endothelin-1 in patients with extra-articular involvement of rheumatoid arthritis. *Clin Rheumatol*. 2006 May;25(3):314-9. DOI: 10.1007/s10067-005-0007-2
- [20] Endemann DH, Schiffrin EL. Endothelial dysfunction. *J Am Soc Nephrol*. 2004 Aug;15(8):1983-92. DOI: 10.1097/01.ASN.0000132474.50966.DA
- [21] Marie I, Bény JL. Endothelial dysfunction in murine model of systemic sclerosis: tight-skin mice 1. *J Invest Dermatol*. 2002 Dec;119(6):1379-87. DOI: 10.1046/j.1523-1747.2002.19614.x
- [22] Strisciuglio T, De Luca S, Capuano E, Luciano R, Niglio T, Trimarco B, Galasso G. Endothelial dysfunction: its clinical value and methods of assessment. *Curr Atheroscler Rep*. 2014 Jun;16(6):417. DOI: 10.1007/s11883-014-0417-1
- [23] Zardi EM, Afeltra A. Endothelial dysfunction and vascular stiffness in systemic lupus erythematosus: Are they early markers of subclinical atherosclerosis? *Autoimmun Rev*. 2010 Aug;9(10):684-6. DOI: 10.1016/j.autrev.2010.05.018
- [24] Ayivi, Raphael & Ibrahim, Salam & Colleran, Heather & Silva, Roberta & Williams, Leonard & Galanakis, Charis & Fidan, Hafize & Tomovska, Julijana & Siddiqui, Shahida. (2021). COVID-19: Human immune response and the influence of food ingredients and active compounds. *Bioactive Compounds in Health and Disease*. 4. 100. DOI: 10.31989/bchd.v4i6.802
- [25] Jourde-Chiche N, Fakhouri F, Dou L, Bellien J, Burtsey S, Frimat M, Jarrot PA, Kaplanski G, Le Quintrec M, Pernin V, Rigotherier C, Sallée M, Fremeaux-Bacchi V, Guerrot D, Roumenina LT. Endothelium structure and function in kidney health and disease. *Nat Rev Nephrol*. 2019 Feb;15(2):87-108. DOI: 10.1038/s41581-018-0098-z
- [26] Yasnaya M.A., Blinov A.V., Blinova A.A., Shevchenko I.M., Maglakelidze D.G., Senkova A.O. Determination of optimal modes for measuring the size of colloidal particles by photon-correlation spectroscopy and acoustic spectroscopy. *Physical and chemical aspects of the study of clusters, nanostructures and nanomaterials*. Tver: TSU, 2020, 12, 232-242. DOI: 10.26456/pcascnn/2020.12.232. (In Russian)
- [27] Cordero MD, de Miguel M, Moreno-Fernández AM. La disfunción mitocondrial en la fibromialgia y su implicación en la patogénesis de la enfermedad [Mitochondrial dysfunction in fibromyalgia and its

implication in the pathogenesis of disease]. *Med Clin (Barc)*. 2011 Mar 12;136(6):252-6. Spanish. DOI: 10.1016/j.medcli.2010.01.030


- [28] Rajendran P, Rengarajan T, Thangavel J, Nishigaki Y, Sakthisekaran D, Sethi G, Nishigaki I. The vascular endothelium and human diseases. *Int J Biol Sci*. 2013 Nov 9;9(10):1057-69. DOI: 10.7150/ijbs.7502
- [29] Kostov K, Halacheva L. Role of Magnesium Deficiency in Promoting Atherosclerosis, Endothelial Dysfunction, and Arterial Stiffening as Risk Factors for Hypertension. *Int J Mol Sci*. 2018 Jun 11;19(6):1724. DOI: 10.3390/ijms19061724.
- [30] Naceiri Mrabti, Hanae & Bouyahya, Abdelhakim & Ed-Dra, Abdelaziz & Kachmar, Mohamed Reda & Naceiri Mrabti, Nidal & Benali, Taoufiq & Shariati, Mohammad Ali & Ouahbi, Abdelilah & Doudach, Latifa & my el abbes, Faouzi. (2021). Polyphenolic profile and biological properties of *Arbutus unedo* root extracts. *European Journal of Integrative Medicine*. 42. 10.1016/j.eujim.2020.101266
- [31] Liguori I, Russo G, Curcio F, Bulli G, Aran L, Della-Morte D, Gargiulo G, Testa G, Cacciatore F, Bonaduce D, Abete P. Oxidative stress, aging, and diseases. *Clin Interv Aging*. 2018 Apr 26;13:757-772. DOI: 10.2147/CIA.S158513
- [32] Daenen K, Andries A, Mekahli D, Van Schepdael A, Jouret F, Bammens B. Oxidative stress in chronic kidney disease. *Pediatr Nephrol*. 2019 Jun;34(6):975-991. DOI: 10.1007/s00467-018-4005-4
-



Ilona Viktorovna Bestaeva is a student of North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Murad Aslambekovich Khasanov is a student of the Medical Institute of the Chechen State University, Grozny, Republic of Chechnya, Russia




Gulshan Anver Babirova is a student of Dagestan State Medical University, Makhachkala, Republic of Dagestan, Russia



Zakhra Zaurovna Askerova, student of Dagestan State Medical University, Makhachkala, Republic of Dagestan, Russia



Dr. Lora Khasanbekovna Dzotsieva, Doctor of Medical Sciences, North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, Russia.



Alina Yurievna Maslova is a Junior Researcher at the Pharmacological Laboratory of Stavropol State Medical University, Stavropol, Russia. Medical Expert, Socmedica, Skolkovo, Moscow, Russia.
